

EXTERIOR BLOCKS AND REFLEXIVE NONCROSSING PARTITIONS

by

Berton Allen Earnshaw

A thesis submitted to the faculty of

Brigham Young University

in partial fulfillment of the requirements for the degree of

Master of Science

Department of Mathematics

Brigham Young University

April 2003

Copyright ©2003 Berton A. Earnshaw  
All Rights Reserved

BRIGHAM YOUNG UNIVERSITY

GRADUATE COMMITTEE APPROVAL

of a thesis submitted by

Berton A. Earnshaw

This thesis has been read by each member of the following graduate committee and by majority vote has been found to be satisfactory.

\_\_\_\_\_  
Date

\_\_\_\_\_  
Rodney W. Forcade, Chair

\_\_\_\_\_  
Date

\_\_\_\_\_  
Lynn E. Garner

\_\_\_\_\_  
Date

\_\_\_\_\_  
Stephen P. Humphries

BRIGHAM YOUNG UNIVERSITY

As chair of the candidate's graduate committee, I have read the thesis of Berton A. Earnshaw in its final form and have found that (1) its format, citations, and bibliographical style are consistent and acceptable and fulfill university and department style requirements; (2) its illustrative materials including figures, tables, and charts are in place; and (3) the final manuscript is satisfactory to the graduate committee and is ready for submission to the university library.

---

Date

---

Rodney W. Forcade  
Chair, Graduate Committee

Accepted for the Department

---

Tyler J. Jarvis  
Graduate Coordinator

Accepted for the College

---

G. Rex Bryce, Associate Dean  
College of Physical and Mathematical Sciences

## ABSTRACT

### EXTERIOR BLOCKS AND REFLEXIVE NONCROSSING PARTITIONS

Berton Allen Earnshaw

Department of Mathematics

Master of Science

This thesis defines an *exterior block* of a noncrossing partition, then gives a formula for the number of noncrossing partitions of the set  $\{1, 2, \dots, n\}$  with  $k$  exterior blocks, which is

$$\frac{k}{n} \binom{2n - k - 1}{k - 1}.$$

Certain identities involving Catalan numbers are derived from this formula. A formula for the number of noncrossing partitions fixed by the reflection of the dihedral group is also derived, which is

$$\binom{n}{\lfloor n/2 \rfloor}$$

the  $n$ th central binomial coefficient.

## ACKNOWLEDGEMENTS

I would like to thank the members of my graduate committee Drs. Rodney W. Forcade, Lynn E. Garner and Stephen P. Humphries for their time, concern and encouragement. I want to especially thank my thesis advisor Dr. Forcade for taking time out of a very busy schedule to help me focus my efforts on this project.

Special thanks is due to Dr. Reinhard O. W. Franz, under whose tutelage I have matured.

I want to thank my wife Tiraje for her love and support, which continues to be the greatest blessing of my life.

# Contents

<b>1</b>	<b>Introduction</b>	<b>8</b>
<b>2</b>	<b>Exterior Blocks</b>	<b>14</b>
<b>3</b>	<b>The Function <math>\text{ext}(n, k)</math></b>	<b>15</b>
<b>4</b>	<b>Catalan Identities</b>	<b>23</b>
<b>5</b>	<b>The Action of the Dihedral Group <math>D_{2n}</math> on the Lattice <math>\text{NC}_n</math></b>	<b>24</b>
<b>6</b>	<b>Reflexive Noncrossing Partitions</b>	<b>26</b>

## List of Figures

1	Linear representation of $1, 4, 6/2, 3/5/7/8, 10/9/11, 12$ . . . . .	9
2	Circular representation of $1, 4, 6/2, 3/5/7/8, 10/9/11, 12$ . . . . .	9
3	Linear representation of $1, 2/3/4, 7/5, 6/8$ . . . . .	11
4	Linear representation of $1, 2, 8/3/4, 7/5, 6$ . . . . .	11
5	Linear representation of $1, 2/3, 8/4, 7/5, 6$ . . . . .	11
6	Linear representation of $1, 2/3/4, 7, 8/5, 6$ . . . . .	12
7	Linear representation of $1, 2/3/4, 7, 8/5, 6$ . . . . .	12
8	Circular representation of $1/2, 4, 7, 9/3/5/6/8$ . . . . .	13
9	Circular representation of $1/2, 4/3/5/6/7, 9/8$ . . . . .	13
10	Circular representation of $1/2, 4/3/5, 6/7, 9/8$ . . . . .	13
11	Circular representation of $1/2, 4, 7, 9/3/5, 6/8$ . . . . .	13
12	Circular representation of $1/2, 4/3, 8/5/6/7, 9$ . . . . .	14
13	Linear representation of $1/2/\cdots/k - 1/k, k + 1, \dots, n$ . . . . .	16
14	Linear representation of $1/\cdots/i - 1/i, i + 1/i + 2/\cdots/n$ . . . . .	17
15	Linear representation of $1, 4, 6/2, 3/5$ . . . . .	17

16	Linear representation of $1, 4/2, 3/5$	18
17	Linear representation of $1, 2/3/4, 5$	18
18	Linear representation of $1, 2, 6/3/4, 5$	18
19	Linear representation of $1, 2/3, 4, 6/5$	19
20	Linear representation of $1, 2/3, 4/5/6$	19
21	Linear representation of $1, 2/3, 4/5, 6$	19
22	Linear representation of $1, 2/3/4, 5, 6$	20
23	Linear representation of $1, 2/3, 6/4, 5$	20
24	Table of values of $\text{ext}(n, k)$	20
25	A Catalan Triangle	22
26	Circular representation of $1, 3/2/4, 5$	25
27	Circular representation of $1, 5/2, 4/3$	25
28	Circular representation of $1, 4/2, 3/5$	26
29	Circular representation of a noncrossing partition	27
30	Circular representation of $2, 3/4$	27
31	Circular representation of $1, 2, 3/4$	28
32	Circular representation of $1, 2, 3, 6, 7/4/5$	28
33	Circular representation of $1, 2, 3, 6, 7/4, 5$	28
34	Circular representation of a noncrossing partition	30
35	Circular representation of $2, 3/4$	32
36	Circular representation of $1, 4, 5/2, 3$	32
37	Circular representation of $1, 4, 5, 6/2, 3/7, 8$	32

## 1 Introduction

In 1972 [13], Kreweras introduced mathematics to *noncrossing partitions*; that is, a partition  $\pi$  of the set  $[n] := \{1, 2, \dots, n\}$  such that whenever  $0 \leq a < b < c < d \leq n$  and  $a$  and  $c$  are in the same block of  $\pi$  and  $b$  and  $d$  are in the same block of  $\pi$ , then ac-



tually  $a$ ,  $b$ ,  $c$  and  $d$  are all in the same block of  $\pi$  (this is the standard definition equivalent, but not equal, to Kreweras' original). The collection of noncrossing partitions of  $[n]$  is denoted by  $\text{NC}_n$ . We typically write noncrossing partitions using a '/' to delimit the blocks of the partition and a ',' to delimit the elements within each block. For example, the partition  $\pi = \{\{1, 4, 6\}, \{2, 3\}, \{5\}, \{7\}, \{8, 10\}, \{9\}, \{11, 12\}\} \in \text{NC}_{12}$  is typically written  $\pi = 1, 4, 6/2, 3/5/7/8, 10/9/11, 12$ . Notice that we have written the blocks in ascending order of their least element. Noncrossing partitions can be conveniently visualized in their *linear* or *circular* representations. For the linear representation, we place  $n$  nodes  $1, 2, \dots, n$  on a line, and indicate that two elements are in the same block by drawing an arc in the upper-half plane connecting the two. For the circular representation, we place  $n$  nodes  $1, 2, \dots, n$  on a circle, and indicate that two elements are in the same block by drawing a line segment in the interior of the circle connecting the two. Figures 1 and 2 give the linear and circular representations, respectively, of  $1, 4, 6/2, 3/5/7/8, 10/9/11, 12$ . Throughout this paper we will make use of both representations.




Figure 1: Linear representation of  $1, 4, 6/2, 3/5/7/8, 10/9/11, 12$




Figure 2: Circular representation of  $1, 4, 6/2, 3/5/7/8, 10/9/11, 12$

Since Kreweras' paper, noncrossing partitions have been studied extensively in a variety of fields. For definitions not given in the discussion that follows, refer to [1] and [17]. It is well-known that  $\text{NC}_n$ , ordered by refinement (that is,  $\pi \leq \sigma$  in  $\text{NC}_n$  if for every block  $B \in \pi$  there exists a block  $C \in \sigma$  such that  $B \subseteq C$ ), forms a graded lattice with rank function  $\text{rk}(\pi) = n - |\pi|$  (of course,  $|\pi|$  denotes the number of blocks of  $\pi$ ), and that  $|\text{NC}_n| = C_n$ , where  $C_n = \frac{1}{n+1} \binom{2n}{n}$  is the  $n$ th *Catalan number* [13]. The number of noncrossing partitions of  $[n]$  of rank  $k$ , which is to ask for the number of noncrossing partitions with  $n - k$  blocks, is  $\frac{1}{n} \binom{n}{n-k} \binom{n}{n-k-1}$  [6].  $\text{NC}_n$  possesses an infimum  $\hat{0} = 1/2/\cdots/n$  and a supremum  $\hat{1} = 1, 2, \dots, n$  and its Möbius function is

$$\mu(\text{NC}_n) = \mu_{\text{NC}_n}(\hat{0}, \hat{1}) = (-1)^{n-1} C_{n-1}$$

[13] [8]. Many chain and multichain enumerations have been formulated, including its zeta polynomial

$$Z_{\text{NC}_n}(m) = \frac{1}{n} \binom{mn}{n-1}$$

[13] [6] [7] [8]. It is known that  $\text{NC}_n$  is rank unimodal, rank symmetric, self-dual and admits a symmetric chain decomposition [15].  $\text{NC}_n$  admits various  $R$ -labelings [4] [8], which have been used to characterize all parking functions, which in turn defines a local action of the symmetric group on  $\text{NC}_n$  [18]. The idea of a noncrossing partition has been generalized [14] and used to study classical reflection groups [3]. Noncrossing partitions are intimately connected with binary trees [12] and meanders [9] [10]. Recently, noncrossing partitions have been used to study stationary stochastic processes with freely independent increments [2].

So what possibly could there be left to study about noncrossing partitions? Consider the “easy” [15] problem of proving by induction that  $|\text{NC}_n| = C_n$ . It seems to require the number of ways the singleton  $\{n+1\}$  can be connected to a noncrossing partition  $\pi \in \text{NC}_n$  to get a noncrossing partition  $\pi' \in \text{NC}_{n+1}$ . Figures 3, 4, 5, 6 and 7 illustrate an example of this problem for  $n = 7$  and  $\pi = 1, 2/3/4, 7/5, 6$ . We

can certainly add the singleton  $\{8\}$  to  $\pi$  to form  $\pi' = \pi \cup \{\{8\}\} = 1, 2/3/4, 7/5, 6/8$  as in Figure 3. We could also add it to the block  $\{1, 2\}$  of  $\pi$  to get  $\pi' = (\pi \setminus \{\{1, 2\}\}) \cup \{\{1, 2, 8\}\} = 1, 2, 8/3/4, 7/5, 6$  as in Figure 4; to the block  $\{3\}$  to get  $\pi' = (\pi \setminus \{\{3\}\}) \cup \{\{3, 8\}\} = 1, 2/3, 8/4, 7/5, 6$  as in Figure 5; and to the block  $\{4, 7\}$  to get  $\pi' = (\pi \setminus \{\{4, 7\}\}) \cup \{\{4, 7, 8\}\} = 1, 2/3/4, 7, 8/5, 6$  as in Figure 6. If we add  $\{8\}$  to the block  $\{5, 6\}$ , we get  $\pi' = (\pi \setminus \{\{5, 6\}\}) \cup \{\{5, 6, 8\}\} = 1, 2/3/4, 7/5, 6, 8$ , which is not a noncrossing partition (see Figure 7).

What is peculiar about the block  $\{5, 6\}$ ? Why does adding the singleton  $\{8\}$  to it form a crossing partition? It is the fact that the block  $\{5, 6\}$  is “nested in” the block  $\{4, 7\}$ . None of the other blocks of  $\pi$  are “nested in” blocks of  $\pi$  in this way. Notice that there are four distinct ways of adding  $\{8\}$  to  $\pi$  to get  $\pi'$ : one for each of the “unnested” blocks of  $\pi$  plus the case of adding  $\{8\}$  as singleton to  $\pi$ .




Figure 3: Linear representation of  $1, 2/3/4, 7/5, 6/8$




Figure 4: Linear representation of  $1, 2, 8/3/4, 7/5, 6$




Figure 5: Linear representation of  $1, 2/3, 8/4, 7/5, 6$

Consider a different problem. The dihedral group  $D_{2n}$  acts on the lattice  $\text{NC}_n$  in a natural way (see Section 5 for details). We think of  $D_{2n}$  as being generated by




Figure 6: Linear representation of  $1, 2/3/4, 7, 8/5, 6$




Figure 7: Linear representation of  $1, 2/3/4, 7, 8/5, 6$

the rotation element  $r = (1, 2, \dots, n)$  and the reflection element  $s = (2, n)(3, n - 1) \cdots (\lceil n/2 \rceil, n - \lceil n/2 \rceil + 2)$ . What is this action like? Given any  $\pi \in \text{NC}_n$ , what is the orbit of  $\pi$  under the action  $D_{2n}$ ? How big is this orbit?

Let us consider the noncrossing partition  $\pi = 1/2, 4, 7, 9/3/5/6/8$ , which is fixed by the action of the reflection  $s$  (see Figure 8; notice that  $\pi$  is symmetric about the dotted line). Notice that  $\pi$  can be constructed by reflecting the noncrossing partition  $\beta = 1/2, 4/3/5$  through the dotted line (see Figure 9) and then joining the block  $\{2, 4\}$  to its reflection  $\{7, 9\}$ . Notice that in a similar way we could decide to connect the block  $\{5\}$  to its reflection (see Figure 10), or connect both the blocks  $\{2, 4\}$  and  $\{5\}$  to their respective reflections (see Figure 11) to get a noncrossing partition fixed by the reflection  $s$  (the reflection of the block  $\{1\}$  of  $\pi$  is simply  $\{1\}$ , so we get nothing new by connecting it to its reflection). However, if we try to connect the block  $\{3\}$  to its reflection, we get a crossing partition (see Figure 12).

Why is the block  $\{3\}$  different from the other blocks of  $\beta$ ? Again, the block  $\{3\}$  is “nested in” the block  $\{2, 4\}$ , while the other blocks of  $\beta$  are not “nested in” any other block of  $\beta$ . Notice that we constructed four distinct noncrossing partitions of  $[9]$ , all fixed by the reflection  $s \in D_{18}$ , from the noncrossing partition  $\beta$ : we had the choice to either connect or not connect the two “unnested” blocks of  $\beta$  with their reflections.




Figure 8: Circular representation of  $1/2, 4, 7, 9/3/5/6/8$




Figure 9: Circular representation of  $1/2, 4/3/5/6/7, 9/8$




Figure 10: Circular representation of  $1/2, 4/3/5, 6/7, 9/8$




Figure 11: Circular representation of  $1/2, 4, 7, 9/3/5, 6/8$




Figure 12: Circular representation of  $1/2, 4/3, 8/5/6/7, 9$

It seems that if we are to count the number of noncrossing partitions of  $[n]$  fixed by the reflection  $s \in D_{2n}$ , or to build up  $\text{NC}_{n+1}$  from  $\text{NC}_n$ , or solve any other problem dependent on the “nestedness” of the blocks of our noncrossing partitions, we will need to define and understand this concept of “nested” blocks.

## 2 Exterior Blocks

For ease of discussion we give a preliminary definition. Given a block  $B \in \pi$ , we will denote the least and greatest elements of  $B$  by  $\text{first}(B)$  and  $\text{last}(B)$ , respectively, and will call them the *first* and *last* elements of  $B$ , respectively.

**Definition 2.1.** Let  $\pi \in \text{NC}_n$ . A block  $B \in \pi$  is an *interior block* of  $\pi$  if there exists a block  $C \in \pi$  such that  $\text{first}(C) < \text{first}(B) \leq \text{last}(B) < \text{last}(C)$ . If  $B$  is not an interior block, then it is an *exterior block* of  $\pi$ .

Intuitively, given a noncrossing partition  $\pi$  of  $[n]$ , an interior block of  $\pi$  is one which is nested inside another block in the linear representation of  $\pi$ . An exterior block of  $\pi$  is one which is not nested in any other block. Consider Figure 1 which is the linear representation of  $\pi = 1, 4, 6/2, 3/5/7/8, 10/9/11, 12 \in \text{NC}_{12}$ . It is easy to see that  $\{2, 3\}$ ,  $\{5\}$  and  $\{9\}$  are the interior blocks of  $\pi$ , while  $\{1, 4, 6\}$ ,  $\{7\}$ ,  $\{8, 10\}$  and  $\{11, 12\}$  are the exterior blocks of  $\pi$ .

We now present a few preliminary results concerning exterior blocks.

**Proposition 2.1.** *Let  $\pi \in \text{NC}_n$ . The blocks of  $\pi$  containing the elements 1 and  $n$  are always exterior blocks.*

*Proof.* Let  $A$  be the block of  $\pi$  containing the element 1 and  $B$  the block containing  $n$ . There is no block of  $\pi$  whose first element is less than  $\text{first}(A) = 1$ , thus, by definition,  $A$  is an exterior block of  $\pi$ . Similarly,  $B$  is an exterior block since there is no block of  $\pi$  whose last element is greater than  $\text{last}(B) = n$ .  $\square$

**Corollary 2.1.** *Every noncrossing partition of  $[n]$  has at least one exterior block.*

*Proof.* This result follows immediately from Proposition 2.1 since every noncrossing partition of  $[n]$  has 1 as an element.  $\square$

**Proposition 2.2.** *Let  $\pi \in \text{NC}_n$ .  $\pi$  has one exterior block if and only if the elements 1 and  $n$  are in the same block.*

*Proof.* ( $\Rightarrow$ ) Suppose  $\pi$  has only one exterior block. If the elements 1 and  $n$  are not contained in the same block, then by Proposition 2.1  $\pi$  has at least two exterior blocks, a contradiction. Therefore 1 and  $n$  must be in the same block.

( $\Leftarrow$ ) Suppose 1 and  $n$  are in the same block  $B$  of  $\pi$ . Then every other block  $A \in \pi$  is an interior block of  $\pi$  since

$$1 = \text{first}(B) < \text{first}(A) \leq \text{last}(A) < \text{last}(B) = n.$$

Therefore,  $B$  is the only exterior block of  $\pi$ .  $\square$

### 3 The Function $\text{ext}(n, k)$

Let  $\text{Ext}_{n,k}$  be the subset of  $\text{NC}_n$  consisting of all noncrossing partitions of  $[n]$  with  $k$  exterior blocks and define

$$\text{ext}(n, k) = |\text{Ext}_{n,k}|$$

so that  $\text{ext}(n, k)$  counts the number of noncrossing partitions of  $[n]$  with  $k$  exterior blocks. What sort of function is  $\text{ext}(n, k)$ ?

**Proposition 3.1.**  $\text{ext}(n, k) = 0$  whenever  $k = 0$  or  $k > n$ .

*Proof.* If  $k = 0$ , we are asking how many noncrossing partitions of  $[n]$  have no exterior blocks. By Corollary 2.1 we know that there are no such noncrossing partitions. Thus  $\text{ext}(n, 0) = 0$ .

Since any partition of  $[n]$  can have at most  $n$  blocks, it can have at most  $n$  exterior blocks. So if  $k > n$ ,  $\text{ext}(n, k) = 0$ . □

**Proposition 3.2.**  $\text{ext}(n, k) > 0$  whenever  $k \in [n]$ .

*Proof.* Given  $k \in [n]$ , the noncrossing partition  $1/2/\cdots/k-1/k, k+1, \dots, n$  has  $k$  exterior blocks (see Figure 13). Therefore  $\text{ext}(n, k) > 0$ . □




Figure 13: Linear representation of  $1/2/\cdots/k-1/k, k+1, \dots, n$

**Proposition 3.3.**  $\text{ext}(n, n) = 1$ .

*Proof.* The only noncrossing partition of  $[n]$  having  $n$  blocks is the infimum  $\widehat{0} = 1/2/\cdots/n$  of the lattice  $\text{NC}_n$ . Notice that each singleton of  $\widehat{0}$  is indeed an exterior block of  $\widehat{0}$ , so  $\text{ext}(n, n) = 1$ . □

**Proposition 3.4.**  $\text{ext}(n, n-1) = n-1$ .

*Proof.* If a noncrossing partition  $\pi$  of  $[n]$  has  $n-1$  exterior blocks, then its blocks must all be singletons except for one block containing two *consecutive* elements; that is,

$$\pi = 1/\cdots/i-1/i, i+1/i+2/\cdots/n$$

for some  $i \in [n-1]$  (see Figure 14). There are as many such noncrossing partitions as there are choices of  $i$ , which number is  $n-1$ . Therefore,  $\text{ext}(n, n-1) = n-1$ . □






Figure 14: Linear representation of  $1/\cdots/i-1/i, i+1/i+2/\cdots/n$

**Theorem 3.1.**  $\text{ext}(n, 1) = C_{n-1}$ , where  $C_n = \frac{1}{n+1} \binom{2n}{n}$  is the  $n$ th Catalan number.

*Proof.* We already know from Proposition 3.3 that  $\text{ext}(1, 1) = 1 = C_0$ . Assume  $n > 1$ . By Proposition 2.2, a noncrossing partition  $\pi$  of  $[n]$  with one exterior block necessarily has 1 and  $n$  in the same block. Call this block  $B$  (see Figure 15, where  $n = 6$ ,  $\pi = 1, 4, 6/2, 3/5$  and  $B = \{1, 4, 6\}$ ). The partition

$$\pi' = (\pi \setminus \{B\}) \cup \{B \setminus \{n\}\}$$

is then a noncrossing partition of  $[n-1]$  ( $\pi'$  is simply  $\pi$  with the element  $n$  removed; see Figure 16). Define a map  $\phi : \text{Ext}_{n,1} \rightarrow \text{NC}_{n-1}$  by the above operation  $\pi \mapsto \pi'$ . The map  $\phi$  is clearly invertible, with inverse map  $\phi^{-1}$  given by

$$\phi^{-1}(\sigma) = (\sigma \setminus \{A\}) \cup \{A \cup \{n\}\}$$

where  $\sigma \in \text{NC}_{n-1}$  and  $A$  is the block of  $\sigma$  containing the element 1 (see Figures 17 and 18, where  $n = 6$ ,  $\sigma = 1, 2/3/4, 5$  and  $A = \{1, 2\}$ ). Therefore  $\phi$  is a bijection, proving

$$\text{ext}(n, 1) = |\text{Ext}_{n,1}| = |\text{NC}_{n-1}| = C_{n-1}.$$

□




Figure 15: Linear representation of  $\pi = 1, 4, 6/2, 3/5 \in \text{Ext}_{6,1}$

So far we have only given the value of  $\text{ext}(n, k)$  for particular values of  $k$ . We now prove a recurrence relation involving  $\text{ext}(n, k)$ .




Figure 16: Linear representation of  $\pi' = \phi(\pi) = 1, 4/2, 3/5 \in \text{Ext}_{5,2}$




Figure 17: Linear representation of  $\sigma = 1, 2/3/4, 5 \in \text{Ext}_{5,3}$

**Theorem 3.2.**  $\text{ext}(n, k) = \text{ext}(n - 1, k - 1) + \text{ext}(n, k + 1)$  for  $n \geq 2$  and  $k \geq 1$ .

*Proof.* Clearly  $\text{ext}(n - 1, k - 1)$  counts the number of noncrossing partitions  $\pi$  of  $\text{Ext}_{n,k}$  having the singleton  $\{n\}$  as a block since  $\pi \setminus \{n\} \in \text{Ext}_{n-1, k-1}$ . Thus we want to show that  $\text{ext}(n, k + 1)$  counts the number of noncrossing partitions of  $\text{Ext}_{n,k}$  that do not have  $\{n\}$  as a block. Let  $\text{Ext}'_{n,k}$  be that set.

If  $k \in [n - 1]$  then by the example in Proposition 3.2 there exists a noncrossing partition with  $k$  exterior blocks whose block containing  $n$  is not a singleton. Thus if  $\text{Ext}'_{n,k}$  is empty, then necessarily  $k \geq n$ . But then  $\text{ext}(n, k + 1) = 0$  by Proposition 3.1 and we are done.

If  $\text{Ext}'_{n,k}$  is not empty, then for any  $\pi \in \text{Ext}'_{n,k}$ , let  $B$  be the block of  $\pi$  containing  $n$  and let

$$\pi' = (\pi \setminus \{B\}) \cup \{B \setminus \{n\}, \{n\}\}$$

(see Figures 19 and 20, where  $n = 6$ ,  $k = 2$ ,  $\pi = 1, 2/3, 4, 6/5$  and  $B = \{3, 4, 6\}$ ). Now  $\pi'$  is a noncrossing partition of  $[n]$  with more than  $k$  exterior blocks. Let  $C$  be the block of  $\pi'$  just to the right of  $B \setminus \{n\}$  in the linear representation of  $\pi'$ ; that is,




Figure 18: Linear representation of  $\phi^{-1}(\sigma) = 1, 2, 6/3/4, 5 \in \text{Ext}_{6,1}$

$\text{last}(B \setminus \{n\}) + 1 = \text{first}(C)$  ( $B \setminus \{6\} = \{3, 4\}$  and  $C = \{5\}$  in Figure 20). Let

$$\pi'' = (\pi \setminus \{C, \{n\}\}) \cup \{C \cup \{n\}\}$$

(see Figure 21). Now  $\pi'' \in \text{Ext}_{n,k+1}$ . Define a map  $\psi : \text{Ext}'_{n,k} \rightarrow \text{Ext}_{n,k+1}$  by the above operation  $\pi \mapsto \pi''$ . The map  $\psi$  is clearly invertible with inverse map  $\psi^{-1}$  given by

$$\psi^{-1}(\sigma) = (\sigma \setminus \{A\}) \cup \{D \cup \{n\}, A \setminus \{n\}\}$$

where  $\sigma \in \text{Ext}_{n,k+1}$  and  $A$  is the block of  $\sigma$  containing  $n$  and  $D$  is the block of  $\sigma$  just to the left of  $A$  in the linear representation of  $\sigma$ ; that is,  $\text{last}(D) + 1 = \text{first}(A)$  (see Figures 22 and 23, where  $n = 6$ ,  $k = 2$ ,  $\sigma = 1, 2/3/4, 5, 6$ ,  $A = \{4, 5, 6\}$  and  $D = \{3\}$ ). Therefore,  $\psi$  is a bijection and

$$|\text{Ext}'_{n,k}| = |\text{Ext}_{n,k+1}| = \text{ext}(n, k + 1).$$

We have proven the desired recurrence. □




Figure 19: Linear representation of  $\pi = 1, 2/3, 4, 6/5 \in \text{Ext}'_{6,2}$




Figure 20: Linear representation of  $\pi' = 1, 2/3, 4/5/6 \in \text{Ext}_{6,4}$



Figure 21: Linear representation of  $\pi'' = \psi(\pi) = 1, 2/3, 4/5, 6 \in \text{Ext}_{6,3}$

This recurrence relations allows us to write out a table of values for  $\text{ext}(n, k)$  (see Figure 24). Notice that the values of the first two columns of this table come




Figure 22: Linear representation of  $\sigma = 1, 2/3/4, 5, 6 \in \text{Ext}_{6,3}$




Figure 23: Linear representation of  $\psi^{-1}(\sigma) = 1, 2/3, 6/4, 5 \in \text{Ext}'_{6,2}$

$n \setminus k$	0	1	2	3	4	5	6	7	8	9	10	Total
1	0	1	0	0	0	0	0	0	0	0	0	1
2	0	1	1	0	0	0	0	0	0	0	0	2
3	0	2	2	1	0	0	0	0	0	0	0	5
4	0	5	5	3	1	0	0	0	0	0	0	14
5	0	14	14	9	4	1	0	0	0	0	0	42
6	0	42	42	28	14	5	1	0	0	0	0	132
7	0	132	132	90	48	20	6	1	0	0	0	429
8	0	429	429	297	165	75	27	7	1	0	0	1430
9	0	1430	1430	1001	572	275	110	35	8	1	0	4862
10	0	4862	4862	3432	2002	1001	429	154	44	9	1	16796

Figure 24: Table of values of  $\text{ext}(n, k)$

from Proposition 3.1 and Theorem 3.1, while the rest of the values come from the recurrence relation written as  $\text{ext}(n, k + 1) = \text{ext}(n, k) - \text{ext}(n - 1, k - 1)$ .

Catalan numbers abound in this table. Notice that the second and third columns (corresponding to  $k = 1$  and  $k = 2$ ) contain Catalan numbers. The first column is, of course, given to us by Theorem 3.1. When  $k = 2$  and  $n \geq 2$ , the recurrence relation plus Proposition 3.1 shows us that

$$\text{ext}(n, 2) = \text{ext}(n, 1) - \text{ext}(n - 1, 0) = C_{n-1} - 0 = C_{n-1}.$$

Notice that the  $n$ th row adds up to  $C_n$ . This is clear since the sets

$$\text{Ext}_{n,1}, \text{Ext}_{n,2}, \dots, \text{Ext}_{n,n}$$

partition  $\text{NC}_n$ ; that is, the sets are pairwise disjoint and  $\text{NC}_n = \cup_{k=1}^n \text{Ext}_{n,k}$ . This fact gives

$$C_n = |\text{NC}_n| = |\cup_{k=1}^n \text{Ext}_{n,k}| = \sum_{k=1}^n |\text{Ext}_{n,k}| = \sum_{k=1}^n \text{ext}(n, k). \quad (3.1)$$

Also notice the strong resemblance of this table with the various formulations of *Catalan's triangle* (cf. [11], also sequences A053121, A008315, etc. in [16]). Figure 25 is a typical Catalan triangle. It is also called a Pascal semi-triangle since if  $w(n, k)$  represents the value in the  $n$ th row and  $k$ th column of this table, then for  $n \geq 1$  and  $k \geq 1$ ,  $w(n, k)$  satisfies the recurrence relation

$$w(n, k) = w(n - 1, k - 1) + w(n - 1, k + 1).$$

Notice that the diagonals  $w(2n, 0), w(2n - 1, 1), \dots, w(n, n)$  of this triangle are the rows  $\text{ext}(n + 1, 1), \text{ext}(n + 1, 2), \dots, \text{ext}(n + 1, n + 1)$  in Figure 24.

We are now ready to give a closed formula for  $\text{ext}(n, k)$ .

**Theorem 3.3.**  $\text{ext}(n, k) = \frac{k}{n} \binom{2n-k-1}{n-1}$ .

$n \setminus k$	0	1	2	3	4	5	6	7	8	9	10	Total
0	1	0	0	0	0	0	0	0	0	0	0	1
1	0	1	0	0	0	0	0	0	0	0	0	1
2	1	0	1	0	0	0	0	0	0	0	0	2
3	0	2	0	1	0	0	0	0	0	0	0	3
4	2	0	3	0	1	0	0	0	0	0	0	6
5	0	5	0	4	0	1	0	0	0	0	0	10
6	5	0	9	0	5	0	1	0	0	0	0	20
7	0	14	0	14	0	6	0	1	0	0	0	35
8	14	0	28	0	20	0	7	0	1	0	0	70
9	0	42	0	48	0	27	0	8	0	1	0	126
10	42	0	90	0	75	0	35	0	9	0	1	252

Figure 25: A Catalan Triangle

*Proof.* Let  $f(n, k) = \frac{k}{n} \binom{2n-k-1}{n-1}$ . Notice that if  $k = 0$  then

$$f(n, 0) = \frac{0}{n} \binom{2n-1}{n-1} = 0 = \text{ext}(n, 0)$$

and if  $k = 1$  then

$$f(n, 1) = \frac{1}{n} \binom{2n-2}{n-1} = C_{n-1} = \text{ext}(n, 1)$$

and if  $k > n$  then  $2n - k - 1 < 2n - n - 1 = n - 1$  so that

$$f(n, k) = \frac{k}{n} \binom{2n-k-1}{n-1} = \frac{k}{n} \cdot 0 = 0 = \text{ext}(n, k)$$

if we follow the convention that  $\binom{a}{b} = 0$  whenever  $b > a$ . Hence  $f(n, k)$  satisfies the initial conditions of Proposition 3.1 and Theorem 3.1. It remains to show that this formula satisfies the recurrence relation

$$f(n, k) = f(n-1, k-1) + f(n, k+1);$$

that is,

$$\begin{aligned} \frac{k}{n} \binom{2n-k-1}{n-1} &= \frac{k-1}{n-1} \binom{2(n-1)-(k-1)-1}{(n-1)-1} + \frac{k+1}{n} \binom{2n-(k+1)-1}{n-1} \\ &= \frac{k-1}{n-1} \binom{2n-k-2}{n-2} + \frac{k+1}{n} \binom{2n-k-2}{n-1} \end{aligned}$$

for  $n \geq 2$  and  $k \geq 1$ . Here we go:

$$\begin{aligned}
f(n-1, k-1) + f(n, k+1) &= \frac{k-1}{n-1} \binom{2n-k-2}{n-2} + \frac{k+1}{n} \binom{2n-k-2}{n-1} \\
&= \frac{k-1}{n-1} \binom{2n-k-2}{n-2} + \frac{k+1}{n} \binom{2n-k-2}{n-1} \\
&= \frac{k-1}{n-1} \cdot \frac{(2n-k-2)!}{(n-2)!(n-k)!} \\
&\quad + \frac{k+1}{n} \cdot \frac{(2n-k-2)!}{(n-1)!(n-k-1)!} \\
&= \frac{n(k-1)(2n-k-2)!}{n!(n-k)!} \\
&\quad + \frac{(n-k)(k+1)(2n-k-2)!}{n!(n-k)!} \\
&= \frac{(nk-n)(2n-k-2)!}{n!(n-k)!} \\
&\quad + \frac{(nk+n-k^2-k)(2n-k-2)!}{n!(n-k)!} \\
&= \frac{(2nk-k^2-k)(2n-k-2)!}{n!(n-k)!} \\
&= \frac{k}{n} \cdot \frac{(2n-k-1)(2n-k-2)!}{(n-1)!(n-k)!} \\
&= \frac{k}{n} \cdot \frac{(2n-k-1)!}{(n-1)!(n-k)!} \\
&= \frac{k}{n} \binom{2n-k-1}{n-1} = f(n, k).
\end{aligned}$$

Therefore,  $\text{ext}(n, k) = f(n, k) = \frac{k}{n} \binom{2n-k-1}{n-1}$ . □

## 4 Catalan Identities

Using the formulation  $\text{ext}(n, k) = \frac{k}{n} \binom{2n-k-1}{n-1}$  of Theorem 3.3, we can derive some identities involving Catalan numbers. The first comes by replacing  $\text{ext}(n, k)$  in Equation 3.1 by this formula:

$$C_n = \sum_{k=1}^n \text{ext}(n, k) = \sum_{k=1}^n \frac{k}{n} \binom{2n-k-1}{n-1}.$$

The second identity comes in response to the question posed at the beginning of this paper regarding a proof by induction of the fact  $|\text{NC}_n| = C_n$  (see Section 1).

There we asked the number of ways the element  $n + 1$  can be added to a noncrossing partition  $\pi$  of  $[n]$  to get a noncrossing partition  $\pi'$  of  $[n + 1]$ , and concluded that if  $\pi$  has  $k$  exterior blocks, then there are  $k + 1$  ways to form the new noncrossing partition  $\pi'$ . Since there are  $\text{ext}(n, k)$  noncrossing partitions of  $[n]$  with  $k$  exterior blocks, there are a total of  $(k + 1)\text{ext}(n, k)$  noncrossing partitions of  $[n + 1]$  gotten in this way. Summing these formulae over the possible number of exterior blocks gives

$$C_{n+1} = |\text{NC}_{n+1}| = \sum_{k=1}^n (k + 1)\text{ext}(n, k) = \sum_{k=1}^n \frac{k(k + 1)}{n} \binom{2n - k - 1}{n - 1}.$$

## 5 The Action of the Dihedral Group $D_{2n}$ on the Lattice $\text{NC}_n$

The dihedral group  $D_{2n}$  acts in a natural way on the lattice  $\text{NC}_n$ . We typically present the dihedral group as  $D_{2n} = \langle r, s \mid r^n, s^2, rsrs \rangle$ , and call the generator  $r$  the *rotation* in  $D_{2n}$  and the generator  $s$  the *reflection* in  $D_{2n}$ . Considered as a subgroup of the symmetric group on  $n$  letters, the generators are typically written  $r = (1, 2, \dots, n)$  and  $s = (2, n)(3, n - 1) \cdots ([n/2], n - [n/2] + 2)$  [5].

A *group action* of a group  $G$  on a set  $A$  is a map  $\cdot : G \times A \rightarrow A$  (where  $\cdot(g, a)$  is typically written  $g \cdot a$ , or even  $ga$ ) such that for all  $g_1, g_2 \in G$  and  $a \in A$  the following two properties hold:

1.  $g_1 \cdot (g_2 \cdot a) = (g_1 g_2) \cdot a$ , and
2.  $1 \cdot a = a$  ( $1$  is the identity element of  $G$ ).

It follows from Property (1) that the action of any group  $G$  on a set  $A$  is determined by the action of the generators of  $G$  on the set  $A$  [5].

The group  $D_{2n}$  acts on the lattice  $\text{NC}_n$  by simply permuting the elements  $1, 2, \dots, n$  of the blocks of any noncrossing partition  $\pi \in \text{NC}_n$ . For instance, if  $n = 5$  and  $\pi = 1, 3/2/4, 5 \in \text{NC}_5$ , then  $r = (1, 2, 3, 4, 5)$ ,  $s = (2, 5)(3, 4)$  and

$$r \cdot \pi = (1, 2, 3, 4, 5) \cdot 1, 3/2/4, 5 = 1, 5/2, 4/3 \in \text{NC}_5$$






Figure 26: Circular representation of  $\pi = 1, 3/2/4, 5$




Figure 27: Circular representation of  $r \cdot \pi = 1, 5/2, 4/3$

and

$$s \cdot \pi = (2, 5)(3, 4) \cdot 1, 3/2/4, 5 = 1, 4/2, 3/5 \in \text{NC}_5.$$

Figures 26, 27 and 28 illustrate these examples (the dotted line represents the axis in which the generator  $s$  reflects  $\pi$ ). It is easily seen that the actions of  $r$  and  $s$  on  $\pi$  do not change the block structure (that is, the number and size of each block, the block adjacencies, etc.) of  $\pi$ ; they simply rotate or reflect it. This fact is true for the actions of  $r$  and  $s$  on any noncrossing partition of  $[n]$ .

An interesting property of the action of  $D_{2n}$  on  $\text{NC}_n$  is that it is rank- and order-preserving. To say that the action is *rank-preserving* means that for any  $\pi \in \text{NC}_n$  and  $d \in D_{2n}$ ,  $\text{rk}(\pi) = \text{rk}(d \cdot \pi)$ . To say that the action is *order-preserving* means that if  $\pi \leq \sigma$  for any  $\sigma \in \text{NC}_n$ , then  $d \cdot \pi \leq d \cdot \sigma$ . This follows immediately from the fact that the action does not change the block structure of a noncrossing partition.

Because the action is rank-preserving and the height of  $\text{NC}_n$  is  $n - 1$  (that is, there




Figure 28: Circular representation of  $s \cdot \pi = 1, 4/2, 3/5$

are  $n$  distinct ranks), when  $n > 1$  this action is not transitive; that is, there is more than one orbit of this action. We are thus led to ask about the number and size of the orbits of this action.

## 6 Reflexive Noncrossing Partitions

In this paper we will only consider the orbits of  $\text{NC}_n$  under the action of the subgroup  $\langle s \rangle$  of  $D_{2n}$ ; that is, we will consider which noncrossing partitions of  $[n]$  are fixed by  $s$ . We will call any noncrossing partition fixed by the reflection  $s$  a *reflexive* noncrossing partition. The number of reflexive noncrossing partitions of  $[n]$  will be denoted by  $s(n)$ .

The discussion will be divided into two cases: one in case  $n$  is odd, and the other in case  $n$  is even. We will begin with the case  $n = 2m + 1$  is odd. If  $n = 1$ , then  $\text{NC}_1$  contains only one noncrossing partition, namely 1, which is clearly reflexive. Assume  $n > 1$ . We will show how to construct the reflexive noncrossing partitions of  $[n]$ .

Since  $n > 1$ , it follows that  $m \geq 1$ , and in the circular representation of any noncrossing partition of  $[n]$  there are  $m$  places labelled  $2, 3, \dots, m + 1$  on one side of the axis of reflection (see Figure 29). In these  $m$  places we can put a noncrossing partition  $\beta$  of  $[2, m + 1] = \{2, 3, \dots, m + 1\}$  with  $k$  exterior blocks (see Figure 30). Now, we add the singleton  $\{1\}$  to  $\beta$  by simply adding the block or by adding it to any one of the exterior blocks of  $\beta$  to get a new noncrossing partition  $\beta'$  of  $[m + 1]$  with




Figure 29: The circular representation of any noncrossing partition of  $[n]$ , where  $n = 2m + 1 > 1$ , has  $m$  places  $2, 3, \dots, m + 1$  on one side of the axis of reflection




Figure 30: Circular representation with  $n = 7$ ,  $m = 3$  and  $\beta = 2, 3/4$

$k' \leq k + 1$  exterior blocks (see Figure 31). We then reflect  $\beta'$  in the axis of reflection to get a reflexive noncrossing partition  $\pi \in \text{NC}_n$  (see Figure 32). We can form more reflexive noncrossing partitions from  $\beta'$  by choosing to connect the exterior blocks of  $\beta'$  not containing 1 with their respective reflections (see Figure 33).

There are  $\text{ext}(m, k)$  ways to choose  $\beta$ . If we do not connect  $\{1\}$  to any of the exterior blocks of  $\beta$ , then  $\beta'$  has  $k' = k + 1$  exterior blocks, so that there are  $2^{k'-1} = 2^k$  ways to construct a reflexive noncrossing partition of  $[n]$  from  $\beta'$ . Hence in this way we can construct  $2^k \text{ext}(m, k)$  reflexive noncrossing partitions of  $[n]$ .

If we connect  $\{1\}$  to the first exterior block of  $\beta$ , then  $\beta'$  has  $k' = k$  exterior blocks, so that there are  $2^{k'-1} = 2^{k-1}$  ways to construct a reflexive noncrossing partition of  $[n]$  from  $\beta'$ . Hence in this way we can construct  $2^{k-1} \text{ext}(m, k)$  reflexive noncrossing partitions of  $[n]$ .




Figure 31: Circular representation with  $n = 7$ ,  $m = 3$ , and  $\beta' = 1, 2, 3/4$




Figure 32: Circular representation of the reflection of  $\beta' = 1, 2, 3/4$  through the axis of reflection, giving  $\pi = 1, 2, 3, 6, 7/4/5$




Figure 33: Circular representation of the reflection of  $\beta' = 1, 2, 3/4$  through the axis of reflection and connecting the block  $\{4\}$  with its reflection  $\{5\}$ , giving  $\pi = 1, 2, 3, 6, 7/4, 5$

In general, if we connect  $\{1\}$  to the  $j$ th exterior block of  $\beta$ ,  $\beta'$  has  $k' = k - j + 1$  exterior blocks, so that there are  $2^{k-j+1-1} = 2^{k-j}$  ways to construct a reflexive noncrossing partition of  $[n]$  from  $\beta'$ . Hence in this way we can construct  $2^{k-j}\text{ext}(m, k)$  reflexive noncrossing partitions of  $[n]$ . Summing the expression  $2^{k-j}\text{ext}(m, k)$  from  $j = 0$  to  $k$  ( $j = 0$  corresponds to the case where  $\{1\}$  is not connected to any of the exterior blocks of  $\beta'$ ) gives

$$\sum_{j=0}^k 2^{k-j}\text{ext}(m, k) = (2^{k+1} - 1)\text{ext}(m, k)$$

ways to construct reflexive noncrossing partitions of  $[n]$  from noncrossing partitions of  $m$  having  $k$  exterior blocks. If we now sum this expression over the possible number of exterior blocks we get a formula for  $s(n)$  when  $n > 1$  is odd:

$$\begin{aligned} s(n) &= \sum_{k=1}^m (2^{k+1} - 1)\text{ext}(m, k) \\ &= \sum_{k=1}^m 2^{k+1}\text{ext}(m, k) - \sum_{k=1}^m \text{ext}(m, k) \\ &= \sum_{k=1}^m \frac{2^{k+1}k}{m} \binom{2m-k-1}{m-1} - C_m \\ &= 4(2m-1)C_{m-1} - C_m \\ &= \binom{2m+1}{m} = \binom{n}{\lfloor n/2 \rfloor} \end{aligned}$$

This is the  $n$ th *central binomial coefficient*. Notice that  $\binom{1}{\lfloor 1 \rfloor} = 1 = s(1)$ , so the formula holds for  $n = 1$  as well.

We now consider the case when  $n = 2m$  is even. If  $n = 2$ , then there are two noncrossing partitions of  $[2]$ , namely  $1, 2$  and  $1/2$ , both of which are clearly reflexive. Assume  $n > 2$ . Then  $m \geq 1$ , and in the circular representation of any noncrossing partition of  $[n]$  there will be  $m - 1$  places labelled  $2, 3, \dots, m$  on one side of the axis of reflection (see Figure 34). If we ignore the place  $m + 1$ , then we are in the case studied above, namely counting the number of reflexive noncrossing partitions of  $[n - 1]$ , of




Figure 34: The circular representation of any noncrossing partition of  $[n]$ , where  $n = 2m > 2$ , has  $m - 1$  places  $2, 3, \dots, m$  on one side of the axis of reflection

which there are

$$s(n - 1) = \binom{n - 1}{\lfloor (n - 1)/2 \rfloor} = \binom{2m - 1}{m - 1}. \quad (6.1)$$

Notice that  $\{m + 1\}$  is a singleton in each of noncrossing partitions counted here.

If we ignore the place 1, then we are again in the same case. As with the case of ignoring the place  $m + 1$ , all the noncrossing partitions that we will count now will have  $\{1\}$  as a singleton block. Notice that we have already counted all those reflexive noncrossing partitions that have both  $\{1\}$  and  $\{m + 1\}$  as singleton blocks. So to avoid double-counting, we must avoid constructing them. If we put a noncrossing partition having  $k$  exterior blocks into the  $m - 1$  places  $2, 3, \dots, m$ , then by the procedure described above we can construct

$$\sum_{j=0}^k 2^{k-j} \text{ext}(m - 1, k) = (2^{k+1} - 1) \text{ext}(m - 1, k)$$

reflexive noncrossing partitions from it. Remember that the index  $j$  represents the position of the exterior block to which we are connecting the block  $\{m + 1\}$ , and that  $j = 0$  represents no connection to the block  $\{m + 1\}$ . This is the case we want to eliminate to avoid double-counting, so we simply subtract  $2^k \text{ext}(m - 1, k)$  from the sum above (which amounts to starting the sum at  $j = 1$ ) to get

$$\sum_{j=1}^k 2^{k-j} \text{ext}(m - 1, k) = (2^k - 1) \text{ext}(m - 1, k)$$

more reflexive noncrossing partitions of  $[n]$ . Summing this number over the possible number of exterior blocks gives

$$\begin{aligned}
\sum_{k=1}^{m-1} (2^k - 1) \text{ext}(m-1, k) &= \sum_{k=1}^{m-1} 2^k \text{ext}(m-1, k) - \sum_{k=1}^{m-1} \text{ext}(m-1, k) \\
&= \sum_{k=1}^{m-1} 2^k \text{ext}(m-1, k) - C_{m-1} \\
&= \frac{1}{2} \left( \sum_{k=1}^{m-1} 2^{k+1} \text{ext}(m-1, k) - C_{m-1} \right) - \frac{1}{2} C_{m-1} \quad (6.2) \\
&= \frac{1}{2} s(n-1) - \frac{1}{2} C_{m-1} \\
&= \frac{1}{2} \binom{n-1}{\lfloor (n-1)/2 \rfloor} - \frac{1}{2} C_{m-1} \\
&= \frac{1}{2} \binom{2m-1}{m-1} - \frac{1}{2} C_{m-1}
\end{aligned}$$

What do we have left to count? We have not considered those reflexive noncrossing partitions which have 1 and  $m+1$  in the same block. How do we count these? If we put a noncrossing partition with  $k$  exterior blocks into the  $m-1$  places (see Figure 35), we have a choice to connect or not connect each of the exterior blocks one at a time to the block  $\{1, m+1\}$  (see Figure 36), and then reflect that in the axis of reflection (see Figure 37). We connect the exterior blocks one at a time to avoid double-counting. This is completely analogous to the situation discussed earlier of building up the noncrossing partitions of  $[m]$  from those of  $[m-1]$ . Thus there are

$$C_m \quad (6.3)$$

such reflexive noncrossing partitions.

Now what is left to count? The only reflexive noncrossing partitions we have not yet counted are those which do not have 1 and  $m+1$  in singleton blocks or in the same block. Again, we begin by putting a noncrossing partition  $\beta$  with  $k$  exterior blocks into the  $m-1$  places. If we add 1 to the first exterior block of  $\beta$ , then we have the choice of adding  $m+1$  to any of the other exterior blocks of  $\beta$ . If we add it to




Figure 35: Circular representation with  $n = 8$ ,  $m = 4$ ,  $k = 2$  and noncrossing partition  $\beta = 2, 3/4$




Figure 36: Circular representation adding the block  $\{4\}$  of  $\beta$  to the block  $\{1, 5\}$  to get  $\beta' = 1, 4, 5/2, 3$




Figure 37: Circular representation reflecting  $\beta'$  in the axis of reflection to get  $\pi = 1, 4, 5, 6/2, 3/7, 8$



the second, then we now have a noncrossing partition  $\beta'$  of  $[m + 1]$  with two exterior blocks. We then reflect  $\beta'$  in the axis of reflection to get a reflexive noncrossing partition  $\pi$  of  $[n]$ . Connecting the exterior blocks of  $\beta'$  to their reflections gives the same noncrossing partition  $\pi$  since both contain either 1 or  $m + 1$ . Thus there are as many of these reflexive noncrossing partitions as there are choices of  $\beta$ , which is  $\text{ext}(m - 1, k)$ .

If we instead add  $m + 1$  to the third exterior block of  $\beta$  to get  $\beta'$  and then reflect  $\beta'$  in the axis of reflection, we get a reflexive noncrossing partition  $\pi$  of  $[n]$ . Notice that  $\beta'$  has three exterior blocks, and if we decide to connect the second exterior block of  $\beta'$  with its reflection we get another reflexive noncrossing partition  $\pi$  of  $[n]$ . Thus there are 2 reflexive noncrossing partitions that can be constructed from  $\beta$  in this way. There are  $\text{ext}(m - 1, k)$  choices for  $\beta$ , so there are  $2\text{ext}(m - 1, k)$  reflexive noncrossing partitions constructed in this way.

In general, if we put a noncrossing partition with  $k$  exterior blocks into the  $m - 1$  places, and if we add 1 to the first exterior block and  $m + 1$  to the  $i$ th exterior block, where  $i > 1$ , we can construct  $2^{i-2}$  reflexive noncrossing partitions of  $[n]$  from it. There are  $\text{ext}(m - 1, k)$  choices for the original partition, so there are  $2^{i-2}\text{ext}(m - 1, k)$  reflexive noncrossing partitions that can be constructed in this way. Adding all these possibilities together, we get

$$\sum_{i=2}^k 2^{i-2}\text{ext}(m - 1, k) = \sum_{i=0}^{k-2} 2^i\text{ext}(m - 1, k)$$

reflexive noncrossing partitions of  $[n]$ .

If we instead add 1 to the  $j$ th exterior block of  $\beta$ , we can only add  $m + 1$  to the  $i$ th exterior block of  $\beta$  if  $i > j$ , else we will have a crossing partition if  $i < j$ , or a noncrossing partition we have already counted if  $i = j$  (since then 1 and  $m + 1$  would be in the same block). As above, there will be  $2^{i-j-1}$  ways to get a reflexive noncrossing partition of  $[n]$  from  $\beta$ , and there are  $\text{ext}(m - 1, k)$  choices for  $\beta$ , so there are  $2^{i-j-1}\text{ext}(m - 1, k)$  reflexive noncrossing partitions of  $[n]$  constructed in this way.

Adding all these possibilities together, we get

$$\sum_{i=j+1}^k 2^{i-j-1} \text{ext}(m-1, k) = \sum_{i=0}^{k-j-1} 2^i \text{ext}(m-1, k)$$

reflexive noncrossing partitions of  $[n]$ . So the number of all of the reflexive noncrossing partitions of  $[n]$  that can be constructed in this way from noncrossing partitions with  $k$  exterior blocks put into the  $m-1$  places is

$$\begin{aligned} \sum_{j=1}^{k-1} \sum_{i=j+1}^{k-j-1} 2^i \text{ext}(m-1, k) &= \sum_{j=0}^{k-2} \sum_{i=0}^{k-j-2} 2^i \text{ext}(m-1, k) \\ &= \sum_{j=0}^{k-2} \sum_{i=0}^j 2^i \text{ext}(m-1, k) \\ &= \sum_{j=0}^{k-2} (2^{j+1} - 1) \text{ext}(m-1, k) \\ &= \left[ \sum_{j=0}^{k-2} 2^{j+1} - \sum_{j=0}^{k-2} 1 \right] \text{ext}(m-1, k) \\ &= [2^k - 2 - (k-1)] \text{ext}(m-1, k) \\ &= (2^k - k - 1) \text{ext}(m-1, k) \end{aligned}$$

If we now sum this expression over the possible number of exterior blocks we get

$$\begin{aligned} \sum_{k=1}^{m-1} \sum_{j=0}^{k-2} \sum_{i=0}^j 2^i \text{ext}(m-1, k) &= \sum_{k=1}^{m-1} (2^k - k - 1) \text{ext}(m-1, k) \\ &= \sum_{k=1}^{m-1} 2^k \text{ext}(m-1, k) - \sum_{k=1}^{m-1} (k+1) \text{ext}(m-1, k) \\ &= \sum_{k=1}^{m-1} 2^k \text{ext}(m-1, k) - C_m \\ &= \frac{1}{2} \sum_{k=1}^{m-1} 2^{k+1} \text{ext}(m-1, k) - \frac{1}{2} C_{m-1} + \frac{1}{2} C_{m-1} - C_m \\ &= \frac{1}{2} s(n-1) + \frac{1}{2} C_{m-1} - C_m \\ &= \frac{1}{2} \binom{n-1}{\lfloor (n-1)/2 \rfloor} + \frac{1}{2} C_{m-1} - C_m \\ &= \frac{1}{2} \binom{2m-1}{m-1} + \frac{1}{2} C_{m-1} - C_m \end{aligned}$$

(6.4)

reflexive noncrossing partitions of  $[n]$  constructed in this way.

Therefore, adding up the numbers in Equations 6.1, 6.2, 6.3 and 6.4, which represent the total number of reflexive noncrossing partitions of  $[n]$  constructed in the four different ways, we get a total of

$$\binom{2m-1}{m-1} + \frac{1}{2} \binom{2m-1}{m-1} - \frac{1}{2} C_{m-1} + C_m + \frac{1}{2} \binom{2m-1}{m-1} + \frac{1}{2} C_{m-1} - C_m = 2 \binom{2m-1}{m-1}$$

reflexive noncrossing partitions of  $[n]$  when  $n = 2m > 2$ . But

$$2 \binom{2m-1}{m-1} = 2 \frac{(2m-1)!}{(m-1)!m!} = \frac{(2m)!}{m!m!} = \binom{2m}{m} = \binom{n}{\lfloor n/2 \rfloor}.$$

Notice also that

$$\binom{2}{\lfloor 2/2 \rfloor} = 2 = s(2)$$

so the formula is valid when  $n = 2$  as well. Therefore, we can write down the formula for the number of reflexive noncrossing partitions of  $[n]$  for all  $n \in \mathbb{N}$ :

$$s(n) = \binom{n}{\lfloor n/2 \rfloor}.$$

This is sequence A001405 in [16]. The first terms of this sequence are

$$1, 2, 3, 6, 10, 20, 35, 70, 126, 252, \dots$$

Notice that these are the entries of last column of Figure 25 beginning with row  $n = 1$ .

## References

- [1] Martin Aigner. *Combinatorial theory*. Springer-Verlag, Berlin, 1979.
- [2] Michael Anshelevich. Free stochastic measures via noncrossing partitions. *Adv. Math.*, 155(1):154–179, 2000.
- [3] Christos A. Athanasiadis. On noncrossing and nonnesting partitions for classical reflection groups. *Electron. J. Combin.*, 5(1):Research Paper 42, 16 pp. (electronic), 1998.

- [4] Anders Björner. Orderings of Coxeter groups. In *Combinatorics and algebra (Boulder, Colo., 1983)*, volume 34 of *Contemp. Math.*, pages 175–195. Amer. Math. Soc., Providence, RI, 1984.
- [5] David S. Dummit and Richard M. Foote. *Abstract Algebra*. John Wiley & Sons Inc., New York, second edition, 1999.
- [6] Paul H. Edelman. Chain enumeration and noncrossing partitions. *Discrete Math.*, 31(2):171–180, 1980.
- [7] Paul H. Edelman. Multichains, noncrossing partitions and trees. *Discrete Math.*, 40(2-3):171–179, 1982.
- [8] Paul H. Edelman and Rodica Simion. Chains in the lattice of noncrossing partitions. *Discrete Math.*, 126(1-3):107–119, 1994.
- [9] Reinhard O. W. Franz. A partial order for the set of meanders. *Ann. Comb.*, 2(1):7–18, 1998.
- [10] Reinhard O. W. Franz and Berton A. Earnshaw. A constructive enumeration of meanders. *Ann. Comb.*, 6(1):7–17, 2002.
- [11] Richard K. Guy. Catwalks, sandsteps and Pascal pyramids. *J. Integer Seq.*, 3(1):Article 00.1.6, 1 HTML document (electronic), 2000.
- [12] Martin Klazar. On trees and noncrossing partitions. *Discrete Appl. Math.*, 82(1-3):263–269, 1998.
- [13] G. Kreweras. Sur les partitions non croisées d’un cycle. *Discrete Math.*, 1(4):333–350, 1972.
- [14] Rodica Simion. Combinatorial statistics on type-B analogues of noncrossing partitions and restricted permutations. *Electron. J. Combin.*, 7(1):Research Paper 9, 27 pp. (electronic), 2000.

- [15] Rodica Simion and Daniel Ullman. On the structure of the lattice of noncrossing partitions. *Discrete Math.*, 98(3):193–206, 1991.
- [16] N. J. A. Sloane. The on-line encyclopedia of integer sequences. *The On-Line Encyclopedia of Integer Sequences*.
- [17] Richard P. Stanley. *Enumerative combinatorics. Vol. 1*. Cambridge University Press, Cambridge, 1997. With a foreword by Gian-Carlo Rota, Corrected reprint of the 1986 original.
- [18] Richard P. Stanley. Parking functions and noncrossing partitions. *Electron. J. Combin.*, 4(2):Research Paper 20, approx. 14 pp. (electronic), 1997. The Wilf Festschrift (Philadelphia, PA, 1996).