

NONCOINCIDENCE OF THE STRICT AND STRONG OPERATOR TOPOLOGIES

JOEL H. SHAPIRO

ABSTRACT. Let E be an infinite-dimensional linear subspace of $C(S)$, the space of bounded continuous functions on a locally compact Hausdorff space S . If μ is a regular Borel measure on S , then each element of E may be regarded as a multiplication operator on $L^p(\mu)$ ($1 \leq p < \infty$). Our main result is that the strong operator topology this identification induces on E is properly weaker than the strict topology. For E the space of bounded analytic functions on a plane region G , and μ Lebesgue measure on G , this answers negatively a question raised by Rubel and Shields in [9]. In addition, our methods provide information about the absolutely p -summing properties of the strict topology on subspaces of $C(S)$, and the bounded weak star topology on conjugate Banach spaces.

1. Introduction. Let $C(S)$ denote the space of bounded, continuous, complex valued functions on a locally compact Hausdorff space S , and let $C_0(S)$ denote those functions in $C(S)$ which vanish at infinity. The *strict topology* β on $C(S)$ is the locally convex topology induced by the seminorms

$$f \rightarrow \|fk\|_{\infty} \quad (f \in C(S)),$$

where k runs through $C_0(S)$ and $\|\cdot\|_{\infty}$ denotes the supremum norm. This topology was introduced in [1] by Buck who derived many of its fundamental properties. In particular [1, Theorems 1 and 2]: β is complete-Hausdorff, and weaker than the norm topology; the norm and strictly bounded subsets of $C(S)$ coincide, and the β -dual of $C(S)$ can be identified with $M(S)$, the space of finite, regular Borel measures on S , where the pairing between the spaces is

$$(1.1) \quad \langle f, \mu \rangle = \int f d\mu \quad (f \in C(S), \mu \in M(S)).$$

Let μ be a (possibly infinite) regular Borel measure on S , as defined in [4, Section 52]. Note that built into this definition is the fact that $\mu(K) < \infty$

Received by the editors February 9, 1972.

AMS 1969 subject classifications. Primary 4601, 4625; Secondary 4630

Key words and phrases. Strict topology, strong operator topology, absolutely p -summing topology, bounded continuous functions.

© American Mathematical Society 1972

for every compact subset K of S [4, p. 223]. For each f in $C(S)$ the equation $M_f g = fg$ ($g \in L^p(\mu)$) defines a bounded linear operator M_f on $L^p(\mu)$. It is not difficult to see from the fact that μ gives finite measure to compact sets that the linear map $f \rightarrow M_f$ is actually an isometry taking $C(S)$ into the space of all bounded linear operators on $L^p(\mu)$. Thus $C(S)$ inherits the *strong operator topology* $\sigma_p = \sigma_p(\mu)$, defined by the seminorms

$$(1.2) \quad f \rightarrow \left\{ \int |fg|^p d\mu \right\}^{1/p} \quad (f \in C(S)),$$

where g runs through $L^p(\mu)$ [3, VI. 1.2, p. 475]. Note that σ_p is locally convex and Hausdorff.

Now $C(S)$ also acts on $C_0(S)$ by multiplication, and in this case the corresponding strong operator topology is the strict topology. In [9, 5.18(c), p. 274], Rubel and Shields asked if $\beta = \sigma_2(\mu)$ on the space $H^\infty(G)$, where μ is two-dimensional Lebesgue measure on G , and G supports nonconstant bounded analytic functions. In this case $H^\infty(G)$ is infinite dimensional [9, Section 2.3], so the question is answered in the negative by the following theorem, which is our main result.

THEOREM 1. *Let E be an infinite-dimensional linear subspace of $C(S)$, and suppose μ is a regular Borel measure on S . Then the strong operator topology $\sigma_p(\mu)$ induced on E by its action on $L^p(\mu)$ is properly weaker than the strict topology.*

The proof of this result occupies §3, and uses the notion of absolutely p -summing locally convex topologies, introduced in the next section. In §4 we comment briefly on the bounded strong operator topology and the bounded weak star topology.

2. Absolutely p -summing topologies. Let τ be a locally convex topology on a real or complex linear space E , and let $E' = E'_\tau$ denote the τ -dual of E (all τ -continuous linear functionals on E). For e' in E' and e in E we will write $\langle e, e' \rangle$ instead of $e'(e)$. A sequence (e_n) in E is called *τ -weakly p -summable* if $\sum |\langle e_n, e' \rangle|^p < \infty$ for all e' in E' , and *τ -absolutely p -summable* if $\sum S(e_n)^p < \infty$ for every τ -continuous seminorm S on E ($1 \leq p < \infty$). If every τ -weakly p -summable sequence is τ -absolutely p -summable, we say τ is *absolutely p -summing*. For example, the weak topology on a Banach space is absolutely p -summing for all p ; but if the space is infinite-dimensional, then the Dvoretzky-Rogers theorem [8, Theorem 8, p. 350] asserts that the norm topology is absolutely p -summing for *no* p ($1 \leq p < \infty$). Note that if τ is not absolutely p -summing, then neither is any stronger locally convex topology on E with the same continuous linear functionals.

The following lemma, which is an easy consequence of the Dvoretzky-Rogers theorem, is the key to our proof of Theorem 1. We note that the same idea has been used in [6, Example 2, p. 417].

LEMMA 1. *Let E be an infinite-dimensional normed space, and let F be a linear subspace of E' which norms E ; that is,*

$$(2.1) \quad \|e\| = \sup\{|\langle e, f \rangle| : f \in F, \|f\| \leq 1\}$$

for each e in E . Let τ denote the topology on E of uniform convergence on (norm) null sequences of F . Then τ is not absolutely p -summing ($1 \leq p < \infty$).

PROOF. Since E is infinite-dimensional it follows from the Dvoretzky-Rogers theorem stated above that there is a sequence (e_n) in E which is weakly, but not absolutely, p -summable for the norm topology; that is, $\sum |\langle e_n, e' \rangle|^p < \infty$ for all e' in E' , but $\sum \|e_n\|^p = \infty$. Since τ is weaker than the norm topology, every τ -continuous linear functional on E is norm continuous; hence (e_n) is τ -weakly p -summable. We claim that (e_n) is not τ -absolutely p -summable. For by (2.1) there exists f_n in F with $\|f_n\| \leq 1$, and

$$|\langle e_n, f_n \rangle| > \|e_n\|/2^{1/p} \quad (n = 1, 2, \dots).$$

Let (a_n) be a sequence of nonnegative numbers such that $\lim a_n = 0$, and $\sum a_n^p \|e_n\|^p = \infty$, and let $g_n = a_n f_n$ ($n = 1, 2, \dots$). Then $\lim \|g_n\| = 0$, so the equation $Se = \sup_n |\langle e, g_n \rangle|$ (e in E) defines a τ -continuous seminorm on E . But

$$\begin{aligned} \sum (Se_n)^p &\geq \sum |\langle e_n, g_n \rangle|^p = \sum a_n^p |\langle e_n, f_n \rangle|^p \\ &\geq \sum a_n^p \|e_n\|^p / 2 = \infty, \end{aligned}$$

so τ is not absolutely p -summing. \square

We will also need a result of J. B. Conway concerning factorization of subsets of $M(S)$. Recall that a subset H of $M(S)$ is called *tight* if for each $\varepsilon > 0$ there exists a compact subset K of S such that $|\mu|(S - K) < \varepsilon$ for each μ in H .

LEMMA 2 [2, THEOREM 2.2, p. 476]. *A bounded subset H of $M(S)$ is tight if and only if there is a bounded subset B of $M(S)$ and a function k in $C_0(S)$ such that $H = kB$.*

Here, of course, $kB = \{kb : b \in B\}$. We can now prove the main result of this section.

PROPOSITION 1. *Let E be an infinite-dimensional linear subspace of $C(S)$. Then the strict topology on E is not absolutely p -summing ($1 \leq p < \infty$).*

PROOF. Since the strict dual of $C(S)$ is $M(S)$, where the spaces are paired by (1.1) [1, Theorem 2], it follows easily that the strict dual E'_β of E may be identified with the quotient space $M(S)/E^\circ$, via the pairing

$$\langle f, \alpha + E^\circ \rangle = \int f d\alpha \quad (f \in E, \alpha \in M(S)),$$

where E° is the annihilator of E in $M(S)$ (see [5, Theorem 14.5, p. 120]). Moreover E'_β is a subspace of E' , the norm dual of E , so it is a normed space.

We will need the fact that for each α in $M(S)$ the norm of the coset $\alpha + E^\circ$ viewed as a linear functional on E coincides with its norm as an element of $M(S)/E^\circ$. To see this, note that each e in E acts by integration as a linear functional on $M(S)$ of norm $\|e\|$, so the pairing (1.1) induces an isometric isomorphism of E into $M(S)'$. Standard Banach space theory now shows that the weak star closure \bar{E} of E in $M(S)'$ is isometrically isomorphic to the dual of $M(S)/\bar{E}^\circ$, where \bar{E}° is the annihilator of \bar{E} in $M(S)$. But $\bar{E}^\circ = E^\circ$, which proves our assertion.

Now the evaluation functionals $(\lambda_s : s \in S)$ defined by

$$(2.2) \quad \lambda_s(e) = e(s) \quad (e \text{ in } E)$$

are strictly continuous and have norm ≤ 1 , so E'_β norms E in the sense of Lemma 1; hence Lemma 1 shows that the topology τ of uniform convergence on norm null sequences in E'_β is not absolutely p -summing. Clearly τ is stronger than the weak topology induced on E by E'_β , so we will be finished if we prove that $\tau \leq \beta$; for then $E'_\tau = E'_\beta$, hence β is not absolutely p -summing since τ is not.

To show that $\tau \leq \beta$, suppose (e'_n) is a norm null sequence in E'_β . By the isometric identification of E'_β with $M(S)/E^\circ$ there exists a sequence (α_n) in $M(S)$ such that $\lim \|\alpha_n\| = 0$, and for each n , $\langle e, e'_n \rangle = \int e d\alpha_n$ ($e \in E$). It is easy to see that (the range of) (α_n) is tight, hence by Lemma 2 there is a bounded sequence (λ_n) in $M(S)$ and a function k in $C_0(S)$ such that $\alpha_n = k\lambda_n$ for all n . Thus for e in E ,

$$\sup_n |\langle e, e'_n \rangle| = \sup_n \left| \int ek d\lambda_n \right| \leq \|ek\|_\infty \sup_n \|\lambda_n\|.$$

Since the left side of this inequality is a typical τ -seminorm, and the right side is a β -continuous seminorm, we have $\tau \leq \beta$. \square

Note that Proposition 1 shows in particular that on any infinite-dimensional linear subspace E of $C(S)$ the strict topology is not nuclear. This fact was first conjectured by Klaus D. Bierstedt for $E = H^\infty(D)$, D the open unit disc (private communication).

3. **Proof of Theorem 1.** For convenience we replace the function g in (1.2) by $|g|^p$. Thus the topology $\sigma_p = \sigma_p(\mu)$ is induced by the seminorms

$$(3.1) \quad S_g(f) = \left\{ \int |f|^p g \, d\mu \right\}^{1/p} \quad (f \in E),$$

where g runs through L^+ , the class of nonnegative μ -integrable functions on S . Since the maximum of two L^+ functions is again in L^+ , we see easily that the sets

$$(3.2) \quad \{f \in C(S) : S_g f \leq 1\} \quad (g \in L^+)$$

form a base for the σ_p -neighborhoods of zero in E .

Now if $g \in L^+$, then it follows from the regularity of μ that $g\mu \in M(S)$. By the argument used in the proof of Lemma 2 [2, Theorem 2.2] with $H = \{g\mu\}$, there exists k in $C_0(S)$ and h in L^+ such that $g = k^p h$. Thus

$$S_g f \leq \|fk\|_\infty \|h\|_1^{1/p} \quad (f \in C(S)),$$

so $\sigma_p \leq \beta$ on $C(S)$.

We complete the proof by showing that $\sigma_p \neq \beta$ on E whenever E is infinite-dimensional. If the strict dual E'_β of E is different from the σ_p -dual, then we are done; so suppose these duals coincide. We claim that in this case σ_p is absolutely p -summing; so again $\sigma_p \neq \beta$, this time by Proposition 1.

Recall that the norm on E'_β is the restriction of the E' norm. Suppose (e_n) is a weakly σ_p (hence β) p -summable sequence in E . Then, as in [7, §1.2.3, p. 22], the set

$$\left\{ \sum_1^N a_n e_n : N = 1, 2, \dots; \sum |a_n|^q \leq 1 \right\},$$

where $p^{-1} + q^{-1} = 1$, is bounded in the weak topology induced on E by E'_β , hence strictly bounded by Mackey's theorem [5, §17.5, p. 155]. Since the strict and norm bounded subsets of E coincide [1, Theorem 1], we have

$$\sup \left| \sum_1^N a_n \langle e_n, e' \rangle \right| < \infty,$$

where the supremum is taken over all positive integers N , all sequences (a_n) in the unit ball of l^q , and all e' in the unit ball of E'_β . From this it follows easily that

$$(3.3) \quad \sup \left\{ \sum_1^\infty |\langle e_n, e' \rangle|^p : e' \in E'_\beta, \|e'\| \leq 1 \right\} < \infty.$$

Now if S is a σ_p -continuous seminorm on E , then S is bounded on a set of

the form (3.2), hence $S \leq S_g$ for some g in L^+ . Taking λ_s as in (2.2) we obtain:

$$\begin{aligned} \sum (S e_n)^p &= \sum (S_\alpha e_n)^p = \sum \int |e_n|^p g \, d\mu \\ &= \int \left(\sum |\langle e_n, \lambda_s \rangle|^p \right) g(s) \, d\mu(s) \leq \|g\|_1 \sup \sum |\langle e_n, e' \rangle|^p < \infty, \end{aligned}$$

where the supremum in the last line is taken over all e' in E'_β with $\|e'\| \leq 1$; a condition satisfied by each λ_s . That the supremum is finite follows from (3.3); hence σ_p is an absolutely p -summing topology on E , and $\sigma_p \neq \beta$. \square

4. The bounded weak star and bounded strong operator topologies. Let E be a subspace of $C(S)$, and let $b\sigma_p = b\sigma_p(\mu)$ denote the *bounded strong operator topology* induced on E by its action on $L^p(\mu)$ (see [3, VI. 9.9, p. 512]); that is, the strongest topology on E agreeing with σ_p on norm bounded sets.

If X is a Banach space, then the *bounded weak star topology* on its dual X' is the strongest topology on X' agreeing with the weak star topology on bounded sets [3, V.3.3, p. 427]. According to the Banach-Dieudonné theorem [3, V.5.4], the bounded weak star topology on X' is just the topology of uniform convergence on null sequences of X . From this and Lemma 1 we get the following result, already noted by Lazar and Retheford for $X=c_0$ [6, Example 2, p. 417].

THEOREM 2. *If X is an infinite-dimensional Banach space, then the bounded weak star topology on X' is not absolutely p -summing. In particular, it is not nuclear.*

In [10, Theorem 2, p. 475] we showed that if E is a linear subspace of $C(S)$ whose unit ball is strictly compact, then E is the dual of the quotient Banach space $M(S)/E^\circ$, and the bounded weak star topology thus induced on E is just the strict topology. This quickly yields the following

THEOREM 3. *Suppose E is a linear subspace of $C(S)$ whose unit ball is strictly compact. Let μ be a regular Borel measure on S . Then $b\sigma_p(\mu) = \beta$.*

PROOF. By [10, Theorem 2] β is the strongest topology on E agreeing on bounded sets with the weak topology induced by $M(S)/E^\circ = E'_\beta$. The proof of Theorem 1 shows that $\sigma_p \leq \beta$, so the unit ball of E is also σ_p -compact. But the topology σ_p is Hausdorff, so $\sigma_p = \beta$ on the unit ball of E , hence on every bounded set (since they are both vector topologies). Thus $b\sigma_p = \beta$. \square

In particular note that if E is $H^\infty(G)$ and μ is Lebesgue measure on G ,

then the hypotheses of Theorem 3 are satisfied. Thus if G supports non-constant bounded analytic functions, then $H^\infty(G)$ is infinite-dimensional; and the strict topology on it is the bounded strong operator topology, but not the strong operator topology.

REFERENCES

1. R. C. Buck, *Bounded continuous functions on a locally compact space*, Michigan Math. J. **5** (1958), 95–104. MR **21** #4350.
2. J. B. Conway, *The strict topology and compactness in the space of measures. II*, Trans. Amer. Math. Soc. **126** (1967), 474–486. MR **34** #6503.
3. N. Dunford and J. T. Schwartz, *Linear operators. I: General theory*, Pure and Appl. Math., vol. 7, Interscience, New York, 1958. MR **22** #8302.
4. P. R. Halmos, *Measure theory*, Van Nostrand, Princeton, N.J., 1950. MR **11**, 504.
5. J. Kelley, I. Namioka et al., *Linear topological spaces*, University Series in Higher Math., Van Nostrand, Princeton, N.J., 1963. MR **29** #3851.
6. A. Lazar and J. Rethford, *Nuclear spaces, Schauder bases, and Choquet simplexes*, Pacific J. Math. **37** (1971), 409–419.
7. A. Pietsch, *Nukleare lokalkonvexe Räume*, Schriftenreihe der Institute für Mathematik bei der Deutschen Akademie der Wissenschaften zu Berlin. Reihe A, Reine Mathematik, Heft 1, Akademie-Verlag, Berlin, 1965. MR **31** #6114.
8. ———, *Absolut p -summierende Abbildungen in normierten Räumen*, Studia Math. **28** (1966/67), 333–353. MR **35** #7162.
9. L. A. Rubel and A. L. Shields, *The space of bounded analytic functions on a region*, Ann. Inst. Fourier (Grenoble) **16** (1966), fasc. 1, 235–277. MR **33** #6440.
10. J. Shapiro, *Weak topologies on subspaces of $C(S)$* , Trans. Amer. Math. Soc. **157** (1971), 471–479.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING,
MICHIGAN 48823